

CAHE’s Guideline Reference Guide: Pelvic girdle pain

Guideline elements	European Guidelines
	<u>European guidelines on the diagnosis and treatment of pelvic girdle pain*</u>
Year of issue/ revision	2004
Etiology, risk factor, population	
Stability of pelvic girdle joint	2, 9, 11, 15
Pregnant women	2, 19, 20, 21
Risk factors	2, 21
Red flag	17
Yellow flag	17
Non-pregnant women	2, 19, 22
Pelvic girdle pain	8
Postpartum	31
Diagnosis	
Diagnosis – clinical tests	3, 23
Reliability of the clinical tests	24
Validity of the clinical tests	26
Diagnostic imaging	3, 26
Conventional radiography	27
Computer tomography	27
Magnetic resonance imaging (MRI)	28
Scintigraphy	28
Pain referral maps	28
Diagnostic injection techniques	29
Outcome measures	30
Prevention	
Prevention	36
Treatment	
Treatment	4
Physical therapy	30

Exercises	31
massage	32
Modified back school classes	32
Water gymnastics	33
Acupuncture	33
Using specially shaped pillows	33
Patient advice	33
Manipulation and mobilisation	34
Pelvic belt	34
Electrotherapy	34
SIJ therapeutic injection therapy and radio frequency denervation	34, 35
External fixation	35
Surgery	35

References*: Vleeming A, Albert H B, van der Helm F C T, Lee D, O’stgaard HC, Stuge B, Sturesson B (2004) A definition of joint stability. In: European guidelines on the diagnosis and treatment of pelvic girdle pain. *Cost Action B13; Low back pain: guidelines for its management*. Working group 4 (this report)